


WS-4 9th advanced

①

Task



$$\text{Range} = u \cdot T_d$$

$$= \sqrt{2} \sqrt{\frac{2h}{g}}$$


$$= \sqrt{\frac{4h}{g}}$$

$$u = \sqrt{u_1^2 + u_2^2 + 2u_1u_2 \cos \theta}$$

$$u = \sqrt{u^2 + u^2 + 2u^2 \cos 90^\circ}$$

$$u = \sqrt{2} u$$

②



$$\tan 45^\circ = \frac{h}{R}$$

$$\Rightarrow h = R$$

$$\Rightarrow h = u \sqrt{\frac{2h}{g}}$$

$$\Rightarrow (\sqrt{h})^2 = u \sqrt{\frac{2h}{g}}$$

$$\Rightarrow u = \sqrt{\frac{gh}{2}}$$

Velocity after t' sec


$$v = \sqrt{u^2 + g^2 t'^2}$$

$$= \sqrt{u^2 + g^2 \left(\sqrt{\frac{2h}{g}}\right)^2}$$


$$= \sqrt{\frac{gh}{2} + 2gh}$$

$$= \sqrt{\frac{5gh}{2}}$$

③



θ , with vertical $\sin(90^\circ - \theta) = \cos \theta = \frac{1}{3}$



$$\sin \theta = \frac{\sqrt{8}}{3}$$

$$\text{From } H = \frac{u^2 \sin^2 \theta}{2g} = \frac{(600)^2 \left(\frac{\sqrt{8}}{3}\right)^2}{2 \times 10}$$

$$\text{From fig } \sin \theta = \frac{u_f}{u_s}$$


$$\sin \theta = \frac{720 \times \frac{5}{18}}{600}$$

$$= \frac{600 \times 600 \times \frac{5}{9}}{600} = 16 \text{ km}$$

$$\sin \theta = \frac{1}{3}$$

(4)

$u_2 = 20\sqrt{3} \text{ m/s}$ $u_1 = 20\sqrt{3} \text{ m/s}$



$S_1 = u_1 t$

$= 20\sqrt{3} t$

From $S = ut + \frac{1}{2} at^2$

$\Rightarrow h = \frac{1}{2} g t^2$

$\Rightarrow h = \frac{1}{2} \times 10 t^2$

$\Rightarrow h = 5(12)^2$

$= 720 \text{ m}$

$R = u \times t$

$= 20\sqrt{3} \times 12$

$= 240\sqrt{3}$

$\tan 30^\circ = \frac{R}{h}$

$\Rightarrow \frac{1}{\sqrt{3}} = \frac{R}{h}$

$\Rightarrow \frac{1}{\sqrt{3}} = \frac{240\sqrt{3} t}{5 t^2}$

$\Rightarrow t = 12 \text{ s}$

$AB = 2R = 2 \times 240\sqrt{3}$

$= 480\sqrt{3} \text{ m}$

(5)

Given


$R_{earth} = R_{planet}$

$\Rightarrow \frac{u_e^2 \sin 2\theta}{g_e} = \frac{u_p^2 \sin 2\theta}{g_p}$

$\Rightarrow \frac{5^2}{10} = \frac{3^2}{g_p}$

$\Rightarrow g_p = \frac{3^2 \times 10}{5^2} = 3.6 \text{ m/s}^2$

(6)



$R = u \sqrt{\frac{2h}{g}}$


$\Rightarrow 10.5 = u \sqrt{\frac{2 \times 1}{10}}$

$\Rightarrow u = 1.5 \sqrt{5} \text{ m/s}$

$R = 5 \times 30 \text{ cm}$

$= 1.5 \text{ m}$

(7)



Given $y = x$

$\Rightarrow u_y t_2 + \frac{1}{2} a_y t_2^2 = u_x t_2 + \frac{1}{2} a_x t_2^2$

$\Rightarrow 0 \times t_2 + \frac{1}{2} g t_2^2 = u t_2 + \frac{1}{2} (0) t_2^2$

$\Rightarrow \frac{1}{2} g t_2^2 = u t_2$

$\Rightarrow t_2 = \frac{2u}{g} = 2 \times \frac{30}{10} = 6 \text{ s}$

$t_2 - t_1 = 6 - 3 = 3 \text{ sec}$

After $t_1 = 3 \text{ s}$

$u_y = u \sin \theta$

$\Rightarrow u_y + a_y t_1 = u$

$\Rightarrow g t_1 = u \Rightarrow t_1 = \frac{u}{g} = \frac{30}{10} = 3$

$$30^\circ = \frac{R}{\frac{3R}{4}}$$

Given $h = 5\text{ m}$

$$R = 10\text{ m}$$

We know $R = u\sqrt{\frac{2h}{g}}$

$$\therefore u\sqrt{\frac{2h}{g}} = 10$$

$$\Rightarrow u\sqrt{\frac{2 \times 5}{10}} = 10$$

$$\Rightarrow u = 10\text{ m/s}$$

(9)

$$u_1 = u \cos 30^\circ \hat{i} + u \sin 30^\circ \hat{j}$$

$$= u \frac{\sqrt{3}}{2} \hat{i} + u \frac{1}{2} \hat{j}$$

$$u_2 = -u \cos 60^\circ \hat{i} + u \sin 60^\circ \hat{j}$$

$$= -\frac{u}{2} \hat{i} + u \frac{\sqrt{3}}{2} \hat{j}$$

$$u_1 - u_2 = \frac{\sqrt{3}u}{2} \hat{i} + \frac{u}{2} \hat{j} + \frac{u}{2} \hat{i} - \frac{\sqrt{3}u}{2} \hat{j}$$


$$= \frac{u}{2} \hat{i} (\sqrt{3} + 1) + \frac{u}{2} (1 - \sqrt{3}) \hat{j}$$

$$|u_1 - u_2| = \sqrt{\left(\frac{u}{2} (\sqrt{3} + 1)\right)^2 + \left(\frac{u}{2} (1 - \sqrt{3})\right)^2}$$

$$= \frac{u}{2} \sqrt{(\sqrt{3} + 1)^2 + (1 - \sqrt{3})^2}$$

$$= \frac{u}{2} \sqrt{2(3 + 1)} = u\sqrt{2}$$

(10)



$$R = \left[\frac{3R}{4}, \frac{R}{4} \right]$$

$$\tan \theta = \frac{y}{x}$$

$$y = x \tan \theta \left(1 - \frac{x}{R} \right)$$

$$\Rightarrow \frac{R}{4} = \frac{3R}{4} \tan \theta \left(1 - \frac{3R}{4R} \right)$$

$$\Rightarrow 1 = 3 \tan \theta \left(1 - \frac{3}{4} \right)$$

$$\Rightarrow \tan \theta = \frac{4}{3}$$

$$\Rightarrow \theta = 53^\circ$$

(14)

Initial velocity = u

$$v = 4u$$

$$\text{From } v^2 - u^2 = 2as$$

$$\Rightarrow (4u)^2 - u^2 = 2gH$$

$$\Rightarrow 15u^2 = 2gH$$

$$\Rightarrow H = \frac{15}{2g} u^2$$

$$R = u \sqrt{\frac{2H}{g}} = u \sqrt{\frac{2 \times 15 u^2}{2g \times g}}$$

$$R = \frac{u^2 \sqrt{15}}{g}$$

8)

Given $u = 600 \text{ kmph}$

$$= 600 \times \frac{5}{18}$$


$$= \frac{500}{3} \text{ m/s}$$

$$\text{Range} = u \sqrt{\frac{2h}{g}} = \frac{500}{3} \sqrt{\frac{2 \times 1960}{9.8}}$$

$$= \frac{500}{3} \sqrt{\frac{2 \times 1960}{9.8}}$$

$$= \frac{500}{3} \times 20 = 333.3 \text{ m}$$

9)



Let n be the no. of steps

$$u = 150 \text{ cm/sec} = 150 \times 10^{-2} \text{ m/s}$$

$$H = 20n \times 10^{-2} \text{ m}; R = 20n \times 10^{-2} \text{ m}$$


$$\text{We know } R = u \sqrt{\frac{2H}{g}}$$

$$20n \times 10^{-2} = 150 \times 10^{-2} \sqrt{\frac{2 \times n \times 20 \times 10^{-2}}{10}}$$

$$\Rightarrow 15 \times 10^{-1} \sqrt{n} = n \Rightarrow n = 3$$

Task

1)



$$v = \sqrt{u^2 + 2gH}$$

$$= \sqrt{(98)^2 + 2(9.8)(0.49)}$$

$$v = 98.98 \text{ m/s}$$

2)


$$u_A = 10 \text{ m/s}; u_B = 20 \text{ m/s}$$

$$t_A = t_B$$

Because time of descent independent on velocity

$$t_d = \sqrt{\frac{2h}{g}}$$

let 'n' be the no. of steps
 'h' be the height of each step
 'x' be the width of each step




$$\text{Range} = u \sqrt{\frac{2h}{g}}$$

$$\Rightarrow nx = u \sqrt{\frac{2h}{g}}$$

$$\Rightarrow nx = u \sqrt{\frac{2nh}{g}} \Rightarrow u = x \cdot n \sqrt{\frac{g}{2nh}}$$

$$\Rightarrow u = x \sqrt{\frac{ng}{2h}}$$

(14)



$$x = u_1 \sqrt{\frac{2h}{g}} = 20 \sqrt{\frac{2 \times 78.04}{9.8}}$$

$$\Rightarrow x = 20 \times 4 = 80$$

$$\text{and } y = u_2 \sqrt{\frac{2h}{g}}$$

$$= 30 \sqrt{\frac{2 \times 78.04}{9.8}}$$


$$= 30 \times 4$$

$$= 120 \text{ m}$$

separation After reaching ground

$$= x + y = 80 + 120 = 200 \text{ m}$$

(15)



$$\tan \theta = \frac{v_y}{v_x} = \frac{gt}{u}$$

$$\Rightarrow \tan 30 = \frac{gt}{u}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{2g}{u} \Rightarrow u = 2\sqrt{3}g$$

(17)

Initial velocity = u

Final velocity = v = 2u

$$\text{From } v = \sqrt{u^2 + 2gh}$$

$$v^2 = u^2 + 2gh$$

$$\Rightarrow 4u^2 = u^2 + 2gh$$

$$\Rightarrow h = \frac{3u^2}{2g}$$

6

Given $u_1 = 20 \text{ m/s}$
 $u_2 = 0$

Given $m_1 = m_2$

According to Law of conservation of Linear momentum

$$P_A = P_B$$


$$\Rightarrow m_1 u_1 + m_2 u_2 = 0$$

$$\Rightarrow m_1 u_1 = -m_2 u_2$$

$$\Rightarrow u_1 = -u_2$$

$$\therefore t = \frac{2\sqrt{u_1 u_2}}{g} = \frac{2\sqrt{20 \times 20}}{10} = \frac{2 \times 20}{10} = 4 \text{ s}$$

8



x, y are ranges

$$x = v \sqrt{\frac{2h}{g}}$$

separation between two bodies


After reaching ground $= x + y$

$$\Rightarrow u_1 \sqrt{\frac{2h}{g}} + u_2 \sqrt{\frac{2h}{g}}$$

$$\Rightarrow v \sqrt{\frac{2h}{g}} + 2v \sqrt{\frac{2h}{g}}$$

$$\Rightarrow x + 2x = 3x$$

9



Given $\frac{m_1}{m_2} = \frac{1}{4}$

$$\Rightarrow \frac{v_2}{v_1} = \frac{1}{4}$$

According to law of conservation of momentum $\Rightarrow v_2 = \frac{v_1}{4}$

$$P_A = P_B$$

$$\Rightarrow m_1 v_1 = -m_2 v_2$$

$$\Rightarrow \frac{v_2}{v_1} = \frac{m_1}{m_2}$$

$$\therefore x = (v_1 + v_2) \sqrt{\frac{2h}{g}}$$

$$= \left[v_1 + \frac{v_1}{4} \right] \sqrt{\frac{2h}{g}}$$

$$\Rightarrow \frac{5v_1}{4} \times \frac{2 \times \frac{v_1}{4}}{g} = \frac{v_1^2 \times 5}{4 \times 10}$$

$$= \frac{v_1^2}{8}$$

(4)

(10)

$$\vec{u} = u_x \hat{i} + u_y \hat{j}$$

$$= u \hat{i} + 0 \hat{j}$$

$$u = u \hat{i}$$

$$u_x = u \cos 0 = u$$

$$u_y = u \sin 0 = 0$$

$$u = v_x \hat{i} + v_y \hat{j}$$

$$= u \hat{i} + g t \hat{j}$$

$$[v_y = u_y + a_y t$$

$$= 0 + g t$$

$$v_y = g t]$$

$$|v - u| = u \hat{i} + g t \hat{j} - u \hat{i} = g t \hat{j}$$

$$= |g t \hat{j}| = g t = g \sqrt{\frac{2h}{g}} = \sqrt{2gh}$$

(15)


$$\text{Given } a = 5 \text{ m/s}^2$$

$$\tan \theta = \frac{a}{g}$$

$$\Rightarrow \tan \theta = \frac{5}{10} = \frac{1}{2}$$

$$\Rightarrow \theta = \tan^{-1}\left(\frac{1}{2}\right)$$

(21)



$$h = 0.2 \text{ m}$$

$$u = 500 \text{ m/s}$$

$$\text{Range} = 100 \text{ m}$$

$$\Rightarrow \frac{h}{5} = \frac{1}{25}$$

$$\text{Range} = 100 \text{ m}$$

$$\Rightarrow u \sqrt{\frac{2h}{g}} = 100$$


$$\Rightarrow 500 \sqrt{\frac{2h}{10}} = 100$$

$$\Rightarrow \sqrt{\frac{h}{5}} = \frac{100}{500} = \frac{1}{5}$$

$$\Rightarrow h = 0.2 \text{ m}$$

$$= 20 \text{ cm}$$


(20)



$$\text{Given } u = 52 \text{ m/s}$$

$$\tan \theta = \frac{5}{12}$$

$$\sin \theta = \frac{5}{13}$$



$$\therefore y = u \sin \theta t - \frac{1}{2} g t^2$$

$$\Rightarrow 15 = 52 \times \frac{5}{13} t - \frac{1}{2} \times 10 t^2$$

$$\Rightarrow 15 = 4 \times 5 t - 5 t^2$$

$$\Rightarrow 3 = 4t - t^2$$

$$\Rightarrow t^2 - 4t + 3 = 0$$

$$\Rightarrow (t-3)(t-1) = 0$$

$$t = 3 \text{ sec or } 1 \text{ sec}$$

$\therefore$ To reach A 1 sec taken

and to reach B 3 sec taken