


8th advanced

⑦

WS-6

- ① In 008 - 11 moving  Here the Velocity of Projection  $u = 3 \text{ m/s}$   
time taken to reach ground = 2s.

Height of bridge

$$h = -ut + \frac{1}{2}gt^2$$

$$h = -(3)(2) + \frac{1}{2} \times 10 \times 2^2$$

$$= -6 + 10 \times 2$$

$$= -6 + 20$$

$$h = 14 \text{ m}$$

- ② Let  $u$  be the velocity with which stone was

Projected (ie) Given  $u = v$

The velocity with which the stone reach ground  $v = 2v$ .

From  $v^2 - u^2 = 2as$  Here  $a = g$ ;  $s = h$

$$\Rightarrow (2v)^2 - v^2 = 2gh$$


$$\Rightarrow 4v^2 - v^2 = 2gh$$

$$\Rightarrow 3v^2 = 2gh$$

$$\Rightarrow h = \frac{3v^2}{2g}$$



③



2 bodies are meet at P.

From  $s = ut + \frac{1}{2}at^2$

| 1st stone (or) body             | 2nd body                                   |
|---------------------------------|--------------------------------------------|
| Freely Falling                  | Vertically Projected                       |
| $u_1 = 0; a_1 = g; s_1 = H - x$ | $u_2 = 100 \text{ m/s}; a_2 = -g; s_2 = x$ |

$$\therefore H - x = u_1 t + \frac{1}{2} g t^2$$

$$\Rightarrow H - x = 0 \cdot t + \frac{1}{2} g t^2$$

$$\Rightarrow H - x = \frac{1}{2} g t^2 \rightarrow \text{①}$$

$$x = u_2 t + \frac{1}{2} (-g) t^2$$

$$\Rightarrow x = 100t - \frac{1}{2} g t^2 \rightarrow \text{②}$$

From ① & ② we get

$$\therefore H - (100t - \frac{1}{2} g t^2) = \frac{1}{2} g t^2$$

$$\Rightarrow 300 - 100t + \frac{1}{2} g t^2 = \frac{1}{2} g t^2$$

$$\Rightarrow 300 - 100t = 0 \Rightarrow 100t = 300$$

$$\Rightarrow t = 3 \text{ sec.}$$

$\therefore$  From

$$\text{① } H - x = \frac{1}{2} g t^2$$

$$\Rightarrow 300 - x = \frac{1}{2} \times 10 \times 3^2$$

$$\Rightarrow 300 - x = 5 \times 9$$


$$\Rightarrow 300 - x = 45 \Rightarrow x = 300 - 45$$

$$\Rightarrow x = 255 \text{ m}$$

Both bodies meet after '3' sec of their start at a height 255m from ground.



④



Let the velocity of projection  $u = 10 \text{ m/s}$   
Height of cliff  $= h = 65 \text{ m}$

The velocity of the stone just before reaching the ground is calculated from  
 $v^2 - u^2 = 2as$

$$\Rightarrow v^2 - 10^2 = 2 \times g \times h$$

$$\Rightarrow v^2 - 100 = 2 \times 10 \times 65$$


$$\Rightarrow v^2 - 100 = 1300$$

$$\Rightarrow v^2 = 1400$$

$$\Rightarrow v = \sqrt{1400} \approx 37 \text{ m/s}$$

⑤

Here Given that helicopter is ascending with a speed  $u = 8 \text{ m/s}$



At the time of chopping the speed of food packet is same as that of helicopter.

$$\therefore \text{From } h = -ut + \frac{1}{2}gt^2$$

$$\Rightarrow 12 = -8t + \frac{1}{2} \times 10 \times t^2$$

$$\Rightarrow 12 = -8t + 5t^2$$

$$\Rightarrow 5t^2 - 8t - 12 = 0$$


Then the roots for 't' can be calculated

$$\text{by using } \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-8) \pm \sqrt{(-8)^2 - 4(5)(-12)}}{2 \times 5}$$

$$t = \frac{8 \pm \sqrt{64 + 2400}}{10} = 2.54 \text{ sec}$$



⑥



let  $H$  = Height reached by a ball from 'O'

$h$  = height of the point 'A' above Top

and the point 'B' below the top of a tower.

'u' be the velocity of projection

From

$$V^2 - u^2 = 2as \rightarrow (1)$$

For 'B'

$$V_B^2 - u^2 = 2gh \quad [\text{Ball is in downward journey}]$$

For 'A'

$$V_A^2 - u^2 = 2(-g)h \quad [\text{Ball is moving up}]$$

$$\Rightarrow V_A^2 - u^2 = -2gh$$

In the question given

$$V_B = 2V_A$$

$$\Rightarrow V_B^2 = 4V_A^2$$

$$\Rightarrow (u^2 + 2gh) = 4(u^2 - 2gh)$$

$$\Rightarrow u^2 + 2gh = 4u^2 - 8gh$$

$$\Rightarrow u^2 - 4u^2 = -2gh - 8gh$$

$$\Rightarrow -3u^2 = -10gh$$

$$\Rightarrow u^2 = \frac{10gh}{3} \rightarrow (2)$$

As the ball moving from O  $\rightarrow$  P then from (1)

$$V_P^2 - u^2 = 2(-g)H$$

but at highest point P

$$V_P = 0$$

$$\Rightarrow 0^2 - u^2 = -2gH$$

$$\Rightarrow u^2 = 2gH$$

$$\Rightarrow 5 \frac{10gh}{3} = 2gH$$

[From 2]


$$\Rightarrow H = \frac{5h}{3}$$





(3)

(7)



let B be the highest point reached by the body.

For the journey A  $\rightarrow$  B:

Initial velocity =  $u$ .

Displacement  $s = 250 - 125$   
 $= 125 \text{ m}$ .

Acceleration  $a = -g = -10 \text{ m/s}^2$

Final velocity  $v_B = 0$

using Relation  $v^2 - u^2 = 2as$

$$\Rightarrow 0^2 - u^2 = 2(-10)(125)$$

$$\Rightarrow -u^2 = -2500$$

$$\Rightarrow u = \sqrt{2500} = 50 \text{ m/s}$$

Now for A  $\rightarrow$  C:

$$u = 50 \text{ m/s}; \quad v_c = v_c$$

$$\text{Displacement} = s_{AC} = -(125 - 70)$$

$$= -55 \text{ m}.$$

$$\text{Acceleration } a = -g = -10 \text{ m/s}^2$$

$$\therefore \text{From } v^2 - u^2 = 2as$$

$$\Rightarrow v_c^2 - (50)^2 = 2(-10)(-55)$$

$$\Rightarrow v_c^2 - 2500 = 1100$$

$$\Rightarrow v_c^2 = 3600 \Rightarrow v_c = \sqrt{3600} = 60 \text{ m/s}$$



8) let the velocity of projection  $u = 19.6 \text{ m/s}$   
Time taken by the body to reach ground  $t = 5 \text{ sec}$

$\therefore$  The displacement of the body when it is  
projected from top of a tower is

$$h = -ut + \frac{1}{2}gt^2$$

$$\Rightarrow h = -19.6(5) + \frac{1}{2}(9.8)(5)^2$$

$$\Rightarrow h = -98.0 + 122.5$$


$$= 24.5 \text{ m}$$

9)



(10)

At the time of dropping



$$V_{\text{body}} = V_{\text{Balloon}} = 12 \text{ m/s}$$

The displacement of the body =

Height from which it is dropped

The height can be calculated by using

$$h = -ut + \frac{1}{2}gt^2$$

$$\Rightarrow 81 = -12t + \frac{1}{2} \times 10 t^2$$

$$\Rightarrow 81 = -12t + 5t^2$$

$$\Rightarrow 5t^2 - 12t - 81 = 0$$

This is in the form of quadratic equation  $ax^2 + bx + c = 0$ 

$$\therefore \text{Roots for } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \text{ Here } x = t$$

$$= \frac{-(-12) \pm \sqrt{(-12)^2 - 4(5)(-81)}}{2 \times 5}$$

$$= \frac{12 \pm \sqrt{144 + 1620}}{10}$$

$$= \frac{12 \pm \sqrt{1764}}{10} = \frac{12 \pm 42}{10}$$

$$= \frac{12+42}{10}$$

$$(or) \frac{12-42}{10}$$

$$= \frac{54}{10}$$

There is no 've' value for time.

$$t = 5.4 \text{ sec}$$



(15)

For (i) (ii)

Here the bag is dropped from helicopter.

So the velocity of bag  $u = 2 \text{ m/s}$ .

Then the separation b/w bag and helicopter after

2 sec  $h = ut - \frac{1}{2}gt^2$

$$= 2 \times 2 - \frac{1}{2}(9.8)2^2$$

$$= 4 - (9.8)2$$

$$= 4 - 19.6 \text{ m}$$

$$= -15.6 \text{ m} \text{ Because the bag is}$$

moving in downward direction its displacement is -ve.

The velocity of bag after 2 sec

$$v = u - gt$$

$$= 2 - (9.8) \times 2$$

$$= 2 - 19.6$$

$$= -17.6 \text{ m/s}$$

(16)



For 1st body  $h = -ut_1 + \frac{1}{2}gt_1^2$

$$\Rightarrow h = -6u + \frac{g}{2}(6)^2$$

$$h = -6u + 18g \rightarrow (1)$$

For 2nd stone  $h = ut_2 + \frac{1}{2}gt_2^2$

$$\Rightarrow h = 2u + \frac{g}{2}(2)^2$$

$$\Rightarrow h = 2u + 2g \rightarrow (2)$$



From ① and ② we get

$$\Rightarrow 2u + 2g = -6u + 18g$$

$$\Rightarrow 8u = 16g$$

$$\Rightarrow u = 2g$$

sub 'u' value in ② we get

$$\Rightarrow h = 2(2g) + 2g$$

$$= 4g + 2g$$

$$= 6g$$

$$h = 60 \text{ m}$$

(17)

Given balloon starts from rest i.e.  $u = 0$

Its acceleration  $a = \frac{g}{8}$ ; After 8 sec from start

The velocity of balloon is  $v = u + at$

$$v = 0 + \frac{g}{8} \times 8$$

$$\Rightarrow v = g \text{ m/s}$$

The height to which balloon was risen is 'h'. From

this height the stone was released from balloon.

$$\therefore h = ut + \frac{1}{2}at^2 \Rightarrow h = 0 \times 8 + \frac{1}{2} \times \frac{g}{8} (8)^2$$

$$\Rightarrow h = \frac{1}{2} \times \frac{g}{8} \times 64$$

$$\Rightarrow h = 4g$$

$$\therefore \text{For stone } h = -ut + \frac{1}{2}gt^2 \quad [\text{Here } u = v]$$

$$\Rightarrow 4g = -gt + \frac{1}{2}gt^2$$



$$\Rightarrow t+4 = -t + \frac{1}{2} t^2$$

$$\Rightarrow t^2 - 2t = 8$$

$$\Rightarrow t^2 - 2t - 8 = 0$$

$$\Rightarrow t^2 - 4t + 2t - 8 = 0$$

$$\Rightarrow (t-4)(t+2) = 0$$

i.e.  $t = 4$  or  $-2$  but No 've' values for time


So correct ans is  $t = 4$  sec.

Task SAG

Q.2 =

(1000 = 1

①



The velocity of the packet = velocity of Balloon at the time of dropping.

The displacement of package = Height from which it was dropped. It can be

calculated by using  $h = -ut + \frac{1}{2} g t^2$

$$\Rightarrow 39.2 = -9.8t + \frac{1}{2} (9.8) t^2$$

$$\Rightarrow 39.2 = -9.8t + 4.9t^2$$

$$\Rightarrow 8 = -2t + t^2$$

$$\Rightarrow t^2 - 2t - 8 = 0$$

$$\Rightarrow t^2 - 4t + 2t - 8 = 0$$

$$\Rightarrow t(t-4) + 2(t-4) = 0$$


$$\Rightarrow (t-4)(t+2) = 0$$

$$(t-4)(t+2) = 0$$

No 've' values for time so 4s is correct



②



Here  $H$  = Maximum height reached by the body from top of tower

$$\Rightarrow H = \frac{u^2}{2g}$$

The displacement of the body =  $h$ , this can be calculated by using  $v^2 - u^2 = 2as$

$$\Rightarrow (7u)^2 - u^2 = 2gh$$

$$\Rightarrow 49u^2 - u^2 = 2gh$$

$$\Rightarrow 48u^2 = 2gh$$

$$\Rightarrow h = \frac{48}{2g} u^2$$

$\therefore$  The total distance travelled by the body

$$= H + H + h$$

$$= \frac{u^2}{2g} + \frac{u^2}{2g} + \frac{48}{2g} u^2$$

$$= \frac{50}{2g} u^2 = 25 \frac{u^2}{g}$$

③

Given initial velocity  $u = 20 \text{ m/s}$

$h = 80 \text{ m}$ ;  $a = g = 10 \text{ m/s}^2$

The speed with which the stone reach the ground is

calculated by using  $v^2 - u^2 = 2as$

$$\Rightarrow v^2 - (20)^2 = 2(10)(80)$$

$$\Rightarrow v^2 - 400 = 1600$$

$$\Rightarrow v^2 = 2000 \Rightarrow v = \sqrt{2000}$$

$$\Rightarrow v = 20\sqrt{5} \text{ m/s}$$



(3)

(4) Here the balloon is moving with uniform velocity

The velocity of stone = 10 m/s;  $h = 75$  m

The displacement of stone is  $h = -ut + \frac{1}{2}gt^2$

$$75 = -10t + \frac{1}{2} \times 10 \times t^2$$

$$15 = -2t + t^2$$

$$t^2 - 2t - 15 = 0$$

$$t^2 - 5t + 3t - 15 = 0$$

$$t(t-5) + 3(t-5) = 0$$

$$(t-5)(t+3) = 0$$

$$t = 5 \text{ sec or } -3 \text{ sec.}$$

No -ve values for time, so time is 5 sec correct.

(5)

The velocity of helicopter = velocity of food packet = 2 m/s

After 2 sec velocity of food packet =  $v = u - gt$

$$= 2 - 9.8 \times 2$$

$$= 2 - 19.6$$

$$= -17.6 \text{ m/s}$$

-ve sign shows the direction of motion is downwards.

(6)

Here given Acceleration of balloon is  $a = \frac{g}{8}$

$$u = 0$$

$\therefore$  The time taken by the stone to reach ground can be

calculated by using  $h = -ut + \frac{1}{2}at^2$

$$\Rightarrow h = -0 \times t + \frac{1}{2} \times \frac{g}{8} t^2 \Rightarrow h = \frac{g}{16} t^2$$

$$\Rightarrow t^2 = \frac{16h}{g} \Rightarrow t = 4\sqrt{\frac{h}{g}}$$






⑦

Here the body was projected up with a velocity

$$u = 19.6 \text{ m/s}$$



The body crosses point of projection

After time of flight

$$T = \frac{2u}{g} = \frac{2 \times 19.6}{9.8} = 4 \text{ sec}$$

⑧

After an object was dropped, the distance


between balloon and object after 2 sec is

$$= \frac{1}{2} g t^2$$

$$= \frac{1}{2} \times 10 \times 2^2$$

$$= 10 \times 2 = 20 \text{ m}$$

⑨



Here  $g = 10 \text{ m/s}^2$

For Vertically Projected body velocity after  $t$  sec is  $v = u + at$

$$\Rightarrow v_1 = u_1 + a_1 t$$

$$\Rightarrow v_1 = 40 - gt \quad \text{--- (1)}$$

For Freely Falling body  $v_2 = u_2 + a_2 t$

$$\Rightarrow v_2 = 0 + gt$$

$$\Rightarrow v_2 = gt \quad \text{--- (2)}$$


$$\text{Given } v_1 = v_2 \Rightarrow gt = 40 - gt$$

$$\Rightarrow 2gt = 40$$

$$\Rightarrow t = \frac{40}{2g} = 2 \text{ sec.}$$



(10)



The velocity of stone = velocity of balloon  
- at the time of dropping

$$u = 10 \text{ m/s}$$

Time taken by the stone to reach ground

$$h = -ut + \frac{1}{2}gt^2$$

$$\Rightarrow 75 = -10t + \frac{1}{2} \times 10 t^2$$

$$\Rightarrow 15 = -2t + t^2$$

$$\Rightarrow 15 = -2t + t^2 \Rightarrow t^2 - 2t - 15 = 0 \quad (1)$$

$$\Rightarrow t^2 - 5t + 3t - 15 = 0$$

$$\Rightarrow t(t-5) + 3(t-5) = 0$$

$$\Rightarrow (t-5)(t+3) = 0 \quad \text{i.e. } t = 5 \text{ (or) } -3\text{sec}$$

No -ve values for time, so  $t = 5\text{sec}$

∴ The height of the balloon from ground when stone

reaches the ground is  $h = \frac{1}{2}gt^2$

$$\Rightarrow h = \frac{1}{2} \times 10 \times 5^2$$

$$\Rightarrow h = 5 \times 25$$

$$h = 125\text{m}$$



(16)

① For a Freely falling body  $u=0$ ;  $a=g$

$\therefore$  The ratio of distance travelled by the body in successive sec is  $= 1:3:5: \dots (2n-1)$

(ii)

Given  $x =$  distance travelled in  $n^{\text{th}}$  sec.

$\therefore$  From (i)  $S_n = u + \frac{g}{2}(2n-1)$

$x = \frac{g}{2}(2n-1) \rightarrow \text{①}$

In  $(n+1)^{\text{th}}$  sec let  $s$  be the distance travelled by a freely falling body.

$\therefore s = u + \frac{g}{2}(2(n+1)-1) [u=0]$

$\Rightarrow s = 0 + \frac{g}{2}(2n+2-1)$

$\Rightarrow s = \frac{g}{2}(2n+1) \rightarrow \text{②}$

$\Rightarrow s = ng + \frac{g}{2}$

From ①  $x = \frac{g}{2}(2n) - \frac{g}{2} \Rightarrow x = ng - \frac{g}{2}$

$\Rightarrow x + \frac{g}{2} = ng$

From ② By 'sub' 'ng' value we get

$\Rightarrow s = x + \frac{g}{2} + \frac{g}{2}$

$\Rightarrow \boxed{s = x + g}$

$\Rightarrow \text{Q.E.D.}$



(ii) let 'x' be the distance travelled by a FFB in  $n^{\text{th}}$  sec

$\therefore$  From  $s = ut + \frac{a}{2}(2n-1) \rightarrow (1)$

Here  $u=0$ ;  $a=g$

$$\Rightarrow x = 0 + \frac{g}{2}(2n-1)$$

$$\Rightarrow x = \frac{g}{2}(2n-1) = \frac{g}{2}(2n) - \frac{g}{2}$$

$$\Rightarrow x = ng - \frac{g}{2} \rightarrow (2) \Rightarrow x + \frac{g}{2} = ng$$

let 's' be distance travelled in  $(n-1)^{\text{th}}$  sec

$\therefore$  From (1)  $s = 0 + \frac{g}{2}[2(n-1)-1]$

$$\Rightarrow s = \frac{g}{2}(2n-2-1)$$

$$\Rightarrow s = \frac{g}{2}(2n-3)$$

$$\Rightarrow s = ng - \frac{3g}{2} \rightarrow (3)$$

sub 'ng' value in (3) we get


$$\Rightarrow s = x + \frac{g}{2} - \frac{3g}{2}$$

$$\Rightarrow -s = x - \frac{2g}{2}$$

$$\Rightarrow \boxed{s = x - g}$$



(17)



2nd stone is clearly FFB

so  $u_2 = 0$  ;  $a_2 = g$ 

After '1' sec velocity

$$v_2 = u_2 + a_2 t$$

$$v_2 = 0 + g(1)$$

$$\Rightarrow v_2 = g = 10 \text{ m/s}$$

After '1' sec 2nd stone is at A,  $s_2$  is distance travelled by it

$$\therefore s_2 = u_2 t + \frac{1}{2} a_2 t^2$$

$$\Rightarrow s_2 = 0 \times t + \frac{1}{2} \times g t^2 = \frac{1}{2} g t^2$$

$$\Rightarrow s_2 = \frac{1}{2} \times 10 \times 1^2 = 5 \text{ m}$$

let the first stone is at B, and it covers a distance

$$\text{of } s_1 \therefore s_1 = u_1 t + \frac{1}{2} a_1 t^2 \quad [a_1 = -g]$$

$$s_1 = 20 \times 1 + \frac{1}{2} (-g) (1)^2$$

$$= 20 - \frac{1}{2} \times 10$$


$$s_1 = 15 \text{ m}$$

$\therefore$  After '1' sec the distance between two stones is AB.

Acc to Pythagorean Theorem


$$AB^2 = 20^2 + 20^2$$

$$AB = \sqrt{20^2 + 20^2} = 20\sqrt{2} \text{ m}$$





18



clearly the body is freely falling body

$$\therefore u = 0; a = g = 10 \text{ m/s}^2$$

in 1 sec the distance travelled by

a.FFB is

$$S = ut + \frac{1}{2}gt^2$$

$$S = 0(1) + \frac{1}{2}(10)(1)^2$$

$$S = 0 + 5 = 5 \text{ m}$$

The velocity of a FFB after 1 sec is

$$v = u + gt$$

$$v = 0 + (10)(1)$$

$$v = 10 \text{ m/s}$$

After 1 sec the body reaches P with a

velocity 10 m/s. From here onwards gravity disappears.

so the body travels further 15m without gravity

From the definition of velocity

$$v = \frac{\text{displacement}}{\text{time}}$$

$$\Rightarrow 10 = \frac{15}{t}$$

$$\Rightarrow t = \frac{15}{10} = 1.5 \text{ sec}$$

