Exercise 5.2

1. Fill in the blanks in the following table, given that a is the first term, d the common difference and a_n , the nth term of the A.P.

	а	d	n	a_n
(<u>i</u>)	7	3	8	
(ii)	-18		10	0
(iii)	••••	-3	18	-5
(iv)	-18.9	2.5		3.6
(v)	3.5	0	105	

Solutions:

(i) Given,

First term, a = 7

Common difference, d = 3

Number of terms, n = 8,

We have to find the nth term, $a_n = ?$

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Putting the values,

$$=>7+(8-1)3$$

$$=>7+(7)$$
 3

$$=>7+21=28$$

Hence, $a_n = 28$

(ii) Given,

First term, a = -18

Common difference, d = ?

Number of terms, n = 10

Nth term, $a_n = 0$

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Putting the values,

$$0 = -18 + (10-1)d$$

$$18 = 9d$$

$$d = 18/9 = 2$$

Hence, common difference, d = 2

(iii)Given,

First term,

a = ?

Common difference, d = -3 Number of terms,

n = 18

Nth term, $a_n = -5$

As we know, for an A.P.,

 $a_n = a + (n-1)d$

Putting the values,

$$-5 = a + (18 - 1)(-3)$$

$$-5 = a+(17)(-3)$$

$$-5 = a - 51$$

$$a = 51 - 5 = 46$$

Hence, a = 46

(iv) Given,

First term, a = -18.9

Common difference, d = 2.5

Number of terms, n = ?

Nth term, $a_n = 3.6$

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Substituting the values,

$$3.6 = -18.9 + (n-1)2.5$$

$$3.6 + 18.9 = (n-1)2.5$$

$$22.5 = (n-1)2.5$$

$$(n-1) = 22.5/2.5$$

$$n - 1 = 9$$

$$n = 10$$

Hence, n = 10

(v)Given,

First term, a = 3.5

Common difference, d = 0Number of terms in this AP = 105

```
Nth term, a_n = ?
```

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Substituting the values,

$$a_n = 3.5 + (105 - 1)0$$

$$a_n = 3.5 + 104 \times 0$$

$$a_n = 3.5$$

Hence, $a_n = 3.5$

2. Choose the correct choice in the following and justify:

(i) 30th term of the A.P: 10,7, 4, ..., is

(ii) 11^{th} term of the A.P. -3, -1/2, ,2 is

Solutions:

(i) Given here,

$$A.P = 10, 7,4,...$$

Therefore,

First term, a = 10

Common difference, $d = a_2 - a_1 = 7 - 10 = -3$

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Substituting the values;

$$a_{30} = 10 + (30 - 1)(-3)$$

$$a_{30} = 10 + (29)(-3)$$

$$a_{30} = 10 - 87 = -77$$

Therefore, the correct answer is option C.

(ii) Given here,

$$A.P = -3, -1/2, ,2$$

...

Therefore,

First term a = -3

Common difference, $d = a_2 - a_1$

$$\Rightarrow$$
 (-1/2) -(-3) \Rightarrow (-1/2) + 3 = 5/2

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Putting the values;

$$a_{11} = 3 + (11 - 1)(5/2)$$

$$a_{11} = 3 + (10)(5/2)$$

$$a_{11} = -3 + 25$$

$$a_{11} = 22$$

So, 11th term is 22, the answer is option B.

3. In the following APs find the missing term in the boxes.

- (i) 2, , 26
- (ii) , 13, , 3
- (iii) 5, \square , $9\frac{1}{2}$
- (iv) -4, , , , , 6
- (v) , 38, , , , , , , -22

Solutions:

(i) For the given AP,

The first and third term are;

$$a = 2$$

$$a_3 = 26$$

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Therefore, putting the values here,

$$a_3 = 2 + (3-1)d$$

$$26 = 2 + 2d$$

$$24 = 2d$$

$$d = 12$$

$$a_2 = a + (2-1)d$$

$$a_2 = a + d$$

 $a_2 = 2 + 12 = 14$

Therefore, 14 is the missing term.

(ii) For the given,

 $a_2 = 13$ and

$$a_4 = 3$$

As we know, for an A.P.,

$$a_n = a + (n-1) d$$

Therefore, putting the values here,

$$a_2 = a + (2 - 1)d$$

$$a_2 = \mathbf{a} + \mathbf{d}$$

$$a_4 = a + (4-1)d$$
, given $a_4 = 3$.

$$3 = a + 3d$$
 (ii)

On subtracting equation (i) from (ii),-10 = 2d

$$d = -5$$

From equation (i), putting the value of d, we get

$$13 = a + (-5)$$

$$a = 18$$

$$a_3 = 18 + (3-1)(-5)$$

$$= 18+2(-5) = 18-10 = 8$$

Therefore, the missing terms are 18 and 8 respectively.

(iii) For the given A.P.,

$$a = 5$$
 and

$$a_4 = 19/2$$

As we know, for an A.P.,

$$a_4 = a + (n-1)d$$

Therefore, putting the values here,

$$a_4 = a + (4-1)d$$

$$19/2 = 5 + 3d$$

$$(19/2) - 5 = 3d$$

$$3d = 9/2$$

$$d = 3/2$$

$$a_2 = a + (2-1)d$$

$$a_2 = 5 + 3/2$$

$$a_2 = 13/2$$

$$a_3 = a + (3-1)d$$

$$a_3 = 5 + 2 \times 3/2$$

$$a_3 = 8$$

Therefore, the missing terms are 13/2 and 8 respectively.

(iv)For the given

A.P.,
$$a = -4$$
 and

$$a_6 = 6$$

As we know, for an A.P.,

$$a_n = a + (n-1) d$$

Therefore, putting the values here,

$$a_6 = a + (6-1)d$$

$$6 = -4 + 5d$$

$$10 = 5d$$

$$d = 2$$

$$a_2 = a + d = -4 + 2 = -2$$

$$a_3 = a+2d = -4+2(2) = 0$$

$$a_4 = a+3d = -4+3(2) = 2$$

$$a_5 = a+4d = -4+4(2) = 4$$

Therefore, the missing terms are -2, 0, 2, and 4 respectively.

(v) For the given A.P.,

$$a_2 = 38$$

$$a_6 = -22$$

As we know, for an A.P.,

$$a_n = a + (n-1)d$$

Therefore, putting the values here,

$$a_2 = a + (2-1)d$$

$$38 = a + d$$
(i)

$$a_6 = a + (6-1)d$$

$$-22 = a+5d$$
(ii)

On subtracting equation (i) from (ii), we get

$$-22 - 38 =$$

$$4d - 60 = 4d$$

$$d = -15$$

Since,
$$a_2 = a + d \Rightarrow a = a_2 - d = 38 - (-15) = 53$$

$$a_3 = a + 2d = 53 + 2 (-15) = 23$$

$$a_4 = a + 3d = 53 + 3 (-15) = 8$$

$$a_5 = a + 4d = 53 + 4(-15) = -7$$

Therefore, the missing terms are 53, 23, 8, and -7 respectively.

4. Which term of the A.P. 3, 8, 13, 18, ...

is 78?

Solution:

Given the A.P. series as 3, 8, 13, 18, ...

First term, a = 3

Common difference, $d = a_2 - a_1 = 8 - 3 = 5$

Let the n^{th} term of given A.P. be 78. Now as

we know, $a_n = a + (n-1)d$

Therefore,

78 = 3 + (n-1)5

$$75 = (n-1)5$$

$$(n-1) = 15$$

$$n = 16$$

Hence, 16th term of this A.P. is 78.

5. Find the number of terms in each of the following A.P.

(i) 7, 13, 19, ..., 205

(ii) 18,
$$15\frac{1}{2}$$
, 13... -47

Solutions:

(i) Given, 7, 13, 19, ..., 205 is the A.P

Therefore,

First term, a = 7

Common difference, $d = a_2 - a_1 = 13 - 7 = 6$.

Let there are n terms in this A.P.

$$a_n = 205$$

As we know, for an A.P.,

$$a_n = a + (n-1) d$$

Therefore,
$$205 = 7 + (n - 1) 6$$

$$198 = (n - 1) 6$$

$$33 = (n - 1)$$

$$n = 34$$

Therefore, this given series has 34 terms in it.

(ii) Given, 18, $15\frac{1}{2}$, 13... -47 is the A.P.

First term, a = 18

Common difference, $d = a_2 - a_1 = 15.5 - 18$ d = (31-36)/2 = -5/2Let there are n terms in this A.P.

 $a_n = 205$

As we know, for an A.P.,

 $a_n = a+(n-1)d$

-47 = 18 + (n-1)(-5/2)

-47-18 = (n-1)(-5/2)

-65 = (n-1)(-5/2)

(n-1) = -130/-5

(n-1) = 26

n = 27

Therefore, this given A.P. has 27 terms in it.

6. Check whether -150 is a term of the A.P. 11, 8, 5, 2, ... Solution:

For the given series, A.P. 11, 8, 5, 2...

First term, a = 11

Common difference, $d = a_2 - a_1 = 8 - 11 = -3$

Let -150 be the n^{th} term of this A.P.

As we know, for an A.P.,

 $a_n = a + (n-1)d$

-150 = 11 + (n-1)(-3)

-150 = 11-3n +3

-164 = -3n

n = 164/3

Clearly, n is not an integer but a fraction.

Therefore, - 150 is not a term of this A.P.

7. Find the 31st term of an A.P. whose 11th term is 38 and the 16th

term is 73.

Solution:

Given that, 11^{th} term, $a_{11} = 38$ and 16^{th} term, $a_{16} = 73$

We know that,

$$a_n = a + (n-1)d$$

$$a_{11} = a + (11-1)d$$

```
38 = a+10d ......(i)
In the same way,
a_{16} = a + (16-1)d
73 = a+15d ......(ii)
On subtracting equation (i) from (ii), we get
35 = 5d
d = 7
From equation (i), we can write,
38 = a+10\times(7)
38 - 70 = a
a = -32
a_{31} = a + (31-1) d
=-32+30(7)
=-32+210
=178
Hence, 31<sup>st</sup> term is 178.
8. An A.P. consists of 50 terms of which 3<sup>rd</sup> term is 12 and the last term is 106. Find
the 29th term.
Solution:
Given that.
3^{\text{rd}} term, a_3 = 12
50th term, a_{50} = 106
a_n = a + (n-1)d
a_3 = a + (3-1)d
12 = a + 2d ......(i)
In the same way,
a_{50} = a + (50 - 1)d
106 = a + 49d ......(ii)
On subtracting equation (i) from (ii), we get
94 = 47d
d = 2 = common difference
From equation (i), we can put d=2,
12 = a + 2(2)
a = 12 - 4 = 8
a_{29} = a + (29 - 1) d
a_{29} = 8 + (28)2
```

```
a_{29} = 8 + 56 = 64
```

Therefore, 29th term is 64.

9. If the 3^{rd} and the 9^{th} terms of an A.P. are 4 and -8 respectively. Which term of this A.P. is zero.

Solution:

Given that, 3^{rd} term, $a_3 = 4$ and 9^{th} term, $a_9 = -8$ We know that,

 $a_n = a + (n-1)d$ Therefore,

On subtracting equation (i) from (ii), we will get here,

$$-12 = 6d$$
$$d = -2$$

From equation (i), we can write,

4 = a + 2(-2)

4 = a - 4

a = 8

Let n^{th} term of this AP be $0....a_n=0$.

 $a_n = a + (n-1)d$

0 = 8 + (n-1)(-2)

0 =

8-2n+2

2n = 10

n = 5

Hence, 5th term of this A.P. is 0.

10.If 17th term of an A.P. exceeds its 10th term by 7. Find the common difference. Solution:

We know that, for an A.P series;

$$a_n = a + (n-1)d$$

$$a_{17} = a + (17 - 1)d$$

 $a_{17} = a + 16d$

In the same way, $a_{10} = a+9d$

As it is given in the question,

$$a_{17} - a_{10} = 7$$

Therefore,

$$(a+16d)-(a+9d)=7$$

7d = 7

$$d = 1$$

Therefore, the common difference is 1.

11. Which term of the A.P. 3, 15, 27, 39,.. will be 132 more than its 54th term? Solution:

Given A.P. is 3, 15, 27, 39, ... first term, a = 3 common difference, $d = a_2 - a_1 = 15 - 3 = 12$

We know that, $a_n = a+(n-1)d$ Therefore, $a_{54} = a+(54-1)d$ $\Rightarrow 3+(53)(12)$ $\Rightarrow 3+636 = 639$ $a_{54} = 639$

We have to find that term of this A.P, which is 132 more than a_{54} , i.e. = $a_{54} + 132$ => 639 + 132 = 771.

Assume, n^{th} term be 771.

$$a_n = a + (n-1)d$$

$$771 = 3 + (n-1)12$$

$$768 = (n-1)12$$

$$(n-1) = 64$$

$$n = 65$$

Therefore, 65th term was 132 more than 54th term.

12.Two APs have the same common difference. The difference between their 100^{th} term is 100, what is the difference between their 1000^{th} terms? Solution:

Let, the first term of two APs be a_1 and a_2 respectively and the common difference of these APs be d.

For the first A.P., we know, $a_n = a + (n-1)d$ Therefore, $a_{100} = a_1 + (100-1)d$ $= a_1 + 99d$ $a_{1000} = a_1 + (1000-1)d$ $a_{1000} = a_1 + 999d$

For second A.P., we know, $a_n = a + (n-1)d$ Therefore, $a_{100} = a_2 + (100-1)d$ $a_{100} = a_2 + 99d$ $a_{1000} = a_2 + (1000-1)d$

 $a_{1000} = a_2 + 999d$

Given that, difference between 100^{th} term of the two APs = 100

Therefore,
$$(a_1+99d) - (a_2+99d) = 100$$

 $a_1 - a_2 = 100$(i)

Difference between 1000th terms of the two Aps $(a_1 + (999)d) - (a_2 + (999)d) = a_1 + 999d - a_2 - 999d = a_1 - a_2$. From equation (i):

This difference, $a_1 - a_2 = 100$ Hence, the difference between 1000^{th} terms of the two A.P. will also be 100.

13. How many three-digit numbers are divisible by 7?

First three-digit number that is divisible by 7 are;

First number = 105Second number = 105+7=112Third number = 112+7=119

Therefore, 105, 112, 119, ...

All are three-digit numbers are divisible by 7 and thus, all these are terms of an A.P. having first term as 105 and common difference as 7.

As we know, the largest possible three-digit number is 999.

When we divide 999 by 7, the remainder will be 5.

Therefore, 999-5 = 994 is the maximum possible three-digit number that is divisible by 7.

```
Now the series is as follows. 105, 112, 119, ..., 994
```

```
Let 994 be the nth term of this A.P. first term, a = 105 common difference, d = 7 a_n = 994 n = ? As we know, a_n = a+(n-1)d 994=105+(n-1)7 889=(n-1)7 (n-1) = 127 n = 128
```

Therefore, 128 three-digit numbers are divisible by 7.

14. How many multiples of 4 lie between 10 and 250? Solution:

The first multiple of 4 that is greater than 10 is 12.

Next multiple will be 16.

Therefore, the series will be 12, 16, 20, 24, ...

All these are divisible by 4 and thus, all these are terms of an A.P. with first term as 12 and common difference as 4.

When we divide 250 by 4, the remainder will be 2. Therefore, 250 - 2 = 248 is divisible by 4....So, last term is 248 in this series. If you find out which term is it in number, you will know the total number of terms.

```
The series is as follows, now;
12, 16, 20, 24, ..., 248
Let 248 be the n^{\text{th}} term of this A.P. first term, a = 12
common difference, d = 4
a_n = 248
```

```
As we know,

a_n = a+(n-1)d

248 = 12+(n-1)\times 4

236/4 = n-1

59 = n-1

n = 60
```

Therefore, there are 60 multiples of 4 between 10 and 250.

15. For what value of n, are the nth terms of two APs 63, 65, 67, and 3, 10, 17, ... equal?

Solution:

Given two APs as; 63, 65, 67,... and 3, 10, 17,....

Taking first AP,

63, 65, 67, ...

First term, a = 63

Common difference, $d = a_2 - a_1 = 65 - 63 = 2$

We know,
$$n^{th}$$
 term of this A.P. = $a_n = a+(n-1)d$
 $a_n = 63+(n-1)2 = 63+2n-2$
 $a_n = 61+2n$ (1)

Taking second AP,

3, 10, 17, ...

First term, a = 3

Common difference, $d = a_2 - a_1 = 10 - 3 = 7$

We know that,

*n*th term of this A.P. = 3+(n-1)7

$$a_n = 3 + 7n - 7$$

$$a_n = 7n - 4....(2)$$

Given, n^{th} term of these A.P.s are equal to each other.

Equating both these equations, we get,

$$61+2n = 7n-4$$

$$61+4=5n$$

$$5n = 65$$

$$n = 13$$

Therefore, 13th terms of both these A.P.s are equal to each other.

16.Determine the A.P. whose third term is 16 and the 7th term exceeds the

5th term by 12.

Solution:

Therefore, A.P. will be 4, 10, 16, 22, ...

17. Find the 20^{th} term from the last term of the A.P. 3, 8, 13, ..., 253.

Solution:

a = 4

Given A.P. is3, 8, 13, ..., 253 Common difference, d= 5.

Therefore, we can write the given AP in reverse order as; 253, ..., 13, 8, 3

Now for the reverse AP, first term, a = 253

and common difference, d = 3-8 = -5n = 20....We have to find 20th term of reverse AP.

Therefore, using nth term formula, we get,

$$a_{20} = a+(20-1)d$$

 $a_{20} = 253+(19)(-5)$
 $a_{20} = 253-95$

$$a_{20} = 158$$

So, 20^{th} term of reverse AP is 158.....that means 20^{th} term from last of actual AP is 158. Therefore, 20^{th} term from the last term of the AP 3, 8, 13, ..., 253.is 158.

18. The sum of 4^{th} and 8^{th} terms of an A.P. is 24 and the sum of the 6^{th} and 10^{th} terms is 44. Find the first three terms of the A.P. Solution:

We know that, the nth term of the AP is;

$$a_n = a + (n-1)d$$

$$a_4 = a + (4-1)d$$

$$a_4 = a + 3d$$

In the same way, we can write,

$$a_8 = a + 7d$$

$$a_6 = a + 5d$$

$$a_{10} = a + 9d$$

Given that, 4^{th} term + 8^{th} term = 24

$$a_4 + a_8 = 24$$

$$(a+3d) + (a+7d) = 24$$

$$2a+10d = 24$$

$$a+5d = 12$$
 (i)

Given, 6^{th} term + 10^{th} term = 44

$$a_6 + a_{10} = 44$$

$$(a +5d) + (a+9d) = 44$$

$$2a+14d = 44$$

$$a+7d = 22$$
(ii)

On subtracting equation (i) from (ii), we get,

$$2d = 22 - 12$$

$$2d = 10$$

$$d = 5$$

From equation (i), we get,

$$a+5d = 12$$

$$a+5(5)=12$$

$$a+25 = 12$$

$$a = -13$$

$$a_2 = a + d = -13 + 5 = -8$$

$$a_3 = a_2 + d = -8 + 5 = -3$$
.

Therefore, the first three terms of this A.P. are -13, -8, and -3.

19.Subba Rao started work in 1995 at an annual salary of Rs 5000 and received an increment of Rs 200 each year. In which year did his income reach Rs 7000?

Solution:

It can be seen from the given question, that the incomes of Subba Rao increases every year by Rs.200 and hence, forms an AP.

```
Therefore, after 1995, the salaries of each year are; 5000, 5200, 5400, ...
Here, first term, a = 5000
and common difference, d = 200

Let after n^{\text{th}} year, his salary be Rs 7000.
Therefore, by the n^{\text{th}} term formula of AP, a_n = a + (n-1) d
7000 = 5000 + (n-1)200
200(n-1) = 2000
(n-1) = 10
n = 11
Therefore, in 11th year, his salary will be Rs 7000.
```

20.Ramkali saved Rs 5 in the first week of a year and then increased her weekly saving by Rs. 1.75. If in the n^{th} week, her weekly savings become Rs 20.75, find n. Solution:

Given that, Ramkali saved Rs.5 in first week and then started saving each week by Rs.1.75. Hence,

First term, a = 5....this is saving in the 1st week.

and common difference, d = 1.75

Also given,

Saving in the nth week, So nth term of the AP, $a_n = 20.75$

Find, n = ?

As we know, by the n^{th} term formula,

 $a_n = a + (n-1)d$

Therefore,

20.75 = 5 + (n-1)1.75

15.75 = (n-1)1.75

(n-1) = 15.75/1.75 = 1575/175

=>63/7=9

=> n - 1 = 9

=> n = 10

Hence, n is

10.

So, 10^{th} week income is Rs. 20.75